

Q-stars in scalar-tensor gravitational theories in extra dimensions

Athanasios Prikas

Physics Department, National Technical University, Zografou Campus,
157 80 Athens, Greece.¹

Abstract

We present Jordan-Brans-Dicke and general scalar-tensor gravitational theory in extra dimensions in an asymptotically flat or anti de Sitter spacetime. We consider a special gravitating, boson field configuration, a q -star, in 3, 4, 5 and 6 dimensions, within the framework of the above gravitational theory and find that the parameters of the stable stars are a few per cent different from the case of General Relativity.

PACS number(s): 11.27.+d, 04.40.-b

Keywords: Non-topological solitons, Scalar-tensor gravitational theories.

¹e-mail: aprikas@central.ntua.gr

1 Introduction - Jordan-Brans-Dicke theory in any dimension

Boson stars are stable configurations of a massive, complex scalar field coupled to gravity, [1, 2]. Self interactions, [3, 4], the coupling of the matter scalar field to a gauge field, [5], or the rotation of the solitonic object, [6], were taken into account. When the non-relativistic theory admits non-topological soliton solutions, the corresponding relativistic generalizations, which are very large field configurations, are the soliton stars, [7, 8, 9]. Q-balls is a special class of non-topological solitons, appearing in Lagrangians with a global $U(1)$ symmetry, [10], a local $U(1)$, [11], or a non-abelian $SU(3)$ or $SO(3)$ symmetry, [12]. Q-stars have been investigated in various models, with one and two scalar fields, [13], non-abelian symmetries, [14], with a scalar and a fermion field, [15] and with a local $U(1)$ symmetry, [16], in asymptotically anti de Sitter spacetime, [17], and in 4 or more dimensions, [18].

Scalar-tensor gravitational theories appeared in the original papers of Brans and Dicke, and Jordan, [19, 20]. The Newtonian constant G is replaced by the inverse mean value of a scalar field, ϕ_{BD} , and the total action contains kinetic terms for the new field times an ω_{BD} quantity, which is regarded as a constant in the original theory. The theory can be generalized by replacing the constant ω_{BD} by a function, usually of the Brans-Dicke (BD) scalar field, [21, 22]. The theory of a scalar field, with quartic self-interactions, coupled to the metric and the BD scalar formulated in [23]. Various properties of such configurations have been studied in a series of papers, [24, 25], where especially the matter of gravitational memory of boson stars in scalar tensor gravity has been analyzed, [26, 27]. Their results generalized in scalar-tensor gravitational theories with ω_{BD} no more a constant, but a function of the BD scalar, [30, 31, 32].

The purpose of the present work is double: We write the BD gravitational theory in D dimensions and apply our results in a realistic case of a scalar field, admitting q-ball type solutions in the absence of gravity. We compare our results with the corresponding ones obtained in General Relativity and we investigate the influence of the spacetime dimensionality in the parameters of the star.

All the field configurations under consideration are spherically symmetric and static, so we use a static, spherically symmetric metric in D dimensions:

$$ds^2 = -e^\nu dt^2 + e^\lambda d\rho^2 + \rho^2 d\Omega_{D-2}^2, \quad (1)$$

with $g_{tt} = -e^\nu$ and $d\Omega_{D-2}^2$ the line element on a $(D-2)$ -dimensional unit sphere. The action is:

$$S = \frac{1}{16\pi} \int d^D x \sqrt{-g_D} \left[\phi_{\text{BD}}(R - 2\Lambda) - \frac{\omega_{\text{BD}} g^{\mu\nu} \partial^\lambda \phi_{\text{BD}} \partial_\lambda \phi_{\text{BD}}}{\phi_{\text{BD}}^2} \right] + S_{\text{matter}} , \quad (2)$$

with S_{matter} the contribution to the total action of the matter fields. Λ stands for the negative, or zero, cosmological constant. The Einstein equations take the form:

$$G_{\mu\nu} = \frac{8\pi}{\phi_{\text{BD}}} T_{\mu\nu} + \frac{\omega_{\text{BD}}}{\phi_{\text{BD}}^2} \left(\partial_\mu \phi_{\text{BD}} \partial_\nu \phi_{\text{BD}} - \frac{1}{2} g_{\mu\nu} \partial^\lambda \phi_{\text{BD}} \partial_\lambda \phi_{\text{BD}} \right) + \frac{1}{\phi_{\text{BD}}} \left(\phi_{\text{BD};\mu;\nu} - g_{\mu\nu} \phi_{\text{BD};\lambda}{}^{;\lambda} \right) - \Lambda g_{\mu\nu} , \quad (3)$$

where $T_\mu{}^\nu = \text{diag}(-\varepsilon, p, \dots, p)$ is the energy momentum tensor with trace $T = \varepsilon + Dp$. The equation of motion for the BD field is:

$$\frac{2\omega_{\text{BD}}}{\phi_{\text{BD}}} \phi_{\text{BD};\lambda}{}^{;\lambda} - \frac{\omega_{\text{BD}} \partial^\lambda \phi_{\text{BD}} \partial_\lambda \phi_{\text{BD}}}{\phi_{\text{BD}}^2} + R - 2\Lambda = 0 . \quad (4)$$

Contracting eq. 3, and substituting the trace in eq. 4 we find the final form for the equation of motion of the BD field:

$$\phi_{\text{BD};\lambda}{}^{;\lambda} = \frac{8\pi T - 2\Lambda \phi_{\text{BD}}}{(D-2)\omega_{\text{BD}} + D - 1} . \quad (5)$$

Equation 5 reveals the dimensional sensitivity of the BD theory. Both T and the ω_{BD} -dependent factor depend on the spacetime dimensionality.

We will now find the asymptotic relation for the BD field. For $p \ll \varepsilon$ and $\Lambda = 0$, the tt component of the Einstein equations gives:

$$G_t{}^t = -\frac{8\pi\varepsilon}{\phi_{\text{BD}}} \frac{(D-2)\omega_{\text{BD}} + D}{(D-2)\omega_{\text{BD}} + D - 1} . \quad (6)$$

The corresponding result of the Einstein gravity is:

$$G_t{}^t = -8\pi G_D \varepsilon . \quad (7)$$

We know that for $\varepsilon \ll$, the limit of the BD theory is the Einstein gravity. So, for localized matter configurations, the right boundary value at infinity for the BD field is:

$$\phi_{\text{BD}} = \frac{1}{G_D} \frac{(D-2)\omega_{\text{BD}} + D}{(D-2)\omega_{\text{BD}} + D - 1} . \quad (8)$$

One can verify that eqs. 6 and 8 give the right result for the well-known 4-dimensional case. We will now apply our results to a certain case of gravitating scalar matter, namely q-stars.




Figure 1: The radius of a q-star as a function of ω_{BD} . The numbers within the figures denote the spacetime dimensionality. For $\omega_{\text{BD}} \simeq 1000$ the results of General Relativity are approximately reproduced. In figures 1-4 and 6-14 we use $\omega = 0.45$, equivalently $A_{\text{sur}} = 0.81$. When decreasing ω_{BD} , the star parameters are a few per cent larger.

2 Q-stars in BD gravitational theory

The Lagrangian resulting from the presence of a matter scalar field is:

$$\mathcal{L}_{\text{matter}} = -(\partial_\mu \phi)^* (\partial^\mu \phi) - U , \quad (9)$$

where ϕ is the matter scalar and U a suitable potential, admitting q-ball type solutions in the absence of gravity. We will now insert the q-soliton ansatz:

$$\phi(\vec{\rho}, t) = \sigma(\rho) e^{-i\omega t} . \quad (10)$$

The equation of motion for the scalar field:

$$\phi_{;\lambda}{}^{;\lambda} - \frac{dU}{d|\phi|^2} \phi = 0 , \quad (11)$$

now takes the form:

$$\sigma'' + \left[\frac{D-2}{\rho} + \frac{1}{2}(\nu' - \lambda') \right] \sigma' + e^\lambda \omega^2 e^{-\nu} \sigma - e^\lambda \frac{dU}{d\sigma^2} \sigma = 0 , \quad (12)$$

where the prime denotes the derivative with respect to ρ . We define:

$$A = e^{-\lambda} , \quad B = e^{-\nu} , \quad (13)$$




Figure 2: The value of the matter scalar field at the center of the q-star as a function of ω_{BD} .




Figure 3: The total energy of a q-star as a function of ω_{BD} .




Figure 4: The particle number of a q-star as a function of ω_{BD} .




Figure 5: The ratio of the star mass to its particle number as a function of the particle number for $\omega_{\text{BD}} = 500$. Small values of the particle number correspond to small stars, whose limiting value is the non-gravitating q-ball.

$$\begin{aligned}
W &\equiv e^{-\nu} \left(\frac{\partial \phi}{\partial t} \right)^* \left(\frac{\partial \phi}{\partial t} \right) = e^{-\nu} \omega^2 \sigma^2 , \\
V &\equiv e^{-\lambda} \left(\frac{\partial \phi}{\partial \rho} \right)^* \left(\frac{\partial \phi}{\partial \rho} \right) = e^{-\lambda} \sigma'^2
\end{aligned} \tag{14}$$

and rescale:

$$\begin{aligned}
\tilde{\rho} &= \rho m , \quad \tilde{\omega} = \omega / m , \quad \tilde{\phi} = \phi / m^{\frac{D-2}{2}} , \\
\tilde{U} &= U / m^D , \quad \tilde{W} = W / m^D , \quad \tilde{V} = V / m^D .
\end{aligned} \tag{15}$$

We also rescale for convenience the BD scalar field:

$$\Phi_{\text{BD}} = \phi_{\text{BD}} G_D \frac{(D-2)\omega_{\text{BD}} + D - 1}{(D-2)\omega_{\text{BD}} + D} . \tag{16}$$

Comparing eqs. 8, 16 we find that $\Phi_{\text{BD}} \rightarrow 1$ for $\rho \rightarrow \infty$ and for localized field configurations.

Gravity becomes important when $R^{D-3} \sim G_D M(R)$ where $M(R)$ is a parameter, dependent on the mass trapped within a sphere of radius R . We define:

$$\epsilon \equiv \sqrt{8\pi G_D m^{D-2}} , \tag{17}$$

which is a very small quantity for m of the order of magnitude of some (hundreds) GeV . We find for the interior of a q-star that $U \sim W \sim m^D$, $V \sim \epsilon^2 m^D$, so the energy resulting from the spatial variation of matter field can be neglected within the star. We also define:

$$\tilde{r} = \epsilon \tilde{\rho} . \tag{18}$$

We use a rescaled potential:

$$\tilde{U} = |\tilde{\phi}|^2 \left(1 - |\tilde{\phi}|^2 + \frac{1}{3} |\tilde{\phi}|^4 \right) = \tilde{\sigma}^2 \left(1 - \tilde{\sigma}^2 + \frac{1}{3} \tilde{\sigma}^4 \right) . \tag{19}$$

This potential admits q-ball type solutions in the absence of gravity with $1 > \omega \geq (\sqrt{U/|\phi|^2})_{\text{min}} = 1/2$. Dropping from now on the tildes and the $O(\epsilon)$ quantities, we find an analytical solution for the scalar field:

$$\sigma = (1 + \omega B^{1/2})^{1/2} , \quad W = B\omega^2(1 + \omega B^{1/2}) , \quad U = \frac{1}{3}(1 + \omega^3 B^{3/2}) . \tag{20}$$

The eigenvalue equation for the frequency ω of the scalar field can be found from the equation of motion of the matter field within the surface. As we know from [13, 14, 15], q-type solitons are characterized by a large interior where the scalar field is approximately constant and a thin surface of width of order of m^{-1} , where the scalar field varies rapidly from the σ value at the inner edge of the surface, to the zero value at the outer edge. The soliton exterior is clearly distinguished from the soliton interior and the surface, because the scalar field vanishes at the exterior. The situation is different in the case of *non-solitonic* boson stars, where the star radius is of order of m^{-1} , the scalar field varies from $\sim M_{\text{Pl}}$ at the center of the soliton to the zero value outside and there is no clear distinction between the star interior and the star surface. In the case of solitonic stars, the soliton radius is uniquely defined by the outer edge of the surface. Dropping the $O(\epsilon)$ quantities from the Lagrange equation within the surface we find:

$$V + W - U = 0 . \quad (21)$$

The above equation holds true only within the surface. At the inner edge of the surface σ' and, consequently, V are zero, in order to match the interior with the surface solution, so $W = U$. Using also eq. 20 we find:

$$\omega = \frac{A_{\text{sur}}^{1/2}}{2} = \frac{B_{\text{sur}}^{-1/2}}{2} , \quad (22)$$

where A_{sur} , B_{sur} denote the value of the metrics at the surface of the star. In the absence of gravity, $A(r) = B(r) = 1$, so $\omega = 1/2$.

We will now turn to the Einstein equations. In D dimensions,

$$\begin{aligned} G_t^t &= \frac{A-1}{2r^2}(D-3)(D-2) + \frac{A'}{2r}(D-2) , \\ G_r^r &= \frac{A-1}{2r^2}(D-3)(D-2) - \frac{AB'}{2Br}(D-2) . \end{aligned} \quad (23)$$

The energy-momentum tensor for the matter scalar field is:

$$T_{\mu\nu} = (\partial_\mu\phi)^*(\partial_\nu\phi) + (\partial_\mu\phi)(\partial_\nu\phi)^* - g_{\mu\nu}[g^{\alpha\beta}(\partial_\alpha\phi)^*(\partial_\beta\phi)] - g_{\mu\nu}U . \quad (24)$$

Using eqs. 10, 13-15 and 20 and dropping the $O(\epsilon)$ quantities we can write

the Einstein equations:

$$G_t^t = \frac{(D-2)\omega_{\text{BD}} + D - 1}{[(D-2)\omega_{\text{BD}} + D]\Phi_{\text{BD}}} \times \left[-W - U - \frac{(D-2)W - DU - 2\Lambda\Phi_{\text{BD}} \frac{(D-2)\omega_{\text{BD}} + D}{(D-2)\omega_{\text{BD}} + D - 1}}{(D-2)\omega_{\text{BD}} + D - 1} \right] - \frac{\omega_{\text{BD}} A \Phi_{\text{BD}}'^2}{2\Phi_{\text{BD}}^2} - \frac{AB' \Phi_{\text{BD}}'}{2\Phi_{\text{BD}} B} , \quad (25)$$

$$G_r^r = \frac{(D-2)\omega_{\text{BD}} + D - 1}{[(D-2)\omega_{\text{BD}} + D]\Phi_{\text{BD}}} \times \left[W - U - \frac{(D-2)W - DU - 2\Lambda\Phi_{\text{BD}} \frac{(D-2)\omega_{\text{BD}} + D}{(D-2)\omega_{\text{BD}} + D - 1}}{(D-2)\omega_{\text{BD}} + D - 1} \right] + \frac{\omega_{\text{BD}} A \Phi_{\text{BD}}'^2}{2\Phi_{\text{BD}}^2} + \frac{A \Phi_{\text{BD}}''}{\Phi_{\text{BD}}} + \frac{A' \Phi_{\text{BD}}'}{2\Phi_{\text{BD}}} , \quad (26)$$

and the equation of motion for the BD field:

$$A \left[\Phi_{\text{BD}}'' + \left(\frac{D-2}{r} + \frac{A'}{2A} - \frac{B'}{2B} \right) \Phi_{\text{BD}}' \right] = \frac{(D-2)W - DU - 2\Lambda\Phi_{\text{BD}} \frac{(D-2)\omega_{\text{BD}} + D}{(D-2)\omega_{\text{BD}} + D - 1}}{(D-2)\omega_{\text{BD}} + D} . \quad (27)$$

We numerically solve the coupled system of eqs. 25-27.

We can find the total mass of the field configuration by the relation:

$$A(\rho) = 1 - \frac{2G_D m_\rho}{\rho^{D-3}} - \frac{2\Lambda\rho^2}{(D-2)(D-1)} , \quad (28)$$

where m_ρ is straightforward connected to the total mass, M_ρ , trapped within a sphere of radius ρ :

$$M_\rho = \frac{D-2}{8\pi} \frac{2\pi^{(D-2)/2}}{\Gamma\left(\frac{D-1}{2}\right)} m_\rho , \quad (29)$$

which, with our rescalings, gives for the total mass, M :

$$M = (D-2) \frac{\pi^{(D-1)/2}}{\Gamma\left(\frac{D-1}{2}\right)} r^{D-3} \left[1 - A(r) - \frac{2\Lambda r^2}{(D-2)(D-1)} \right] , \quad r \rightarrow \infty . \quad (30)$$

In the case of a rotating star, one should embed General Relativity in a more general framework, Metric-affine Gravity, where the correct renormalized mass is (in 4 dimensions):

$$\frac{M}{1 + \frac{\Lambda}{3} \frac{J^2}{M^2}} ,$$

with J the total angular momentum of the star, [28, 29]. In our case the anti de Sitter mass is identical with the above renormalized mass, because $J = 0$. The mass of eqs. 28-30 is the Schwarzschild mass, which corresponds to the ADM mass in the Jordan frame. In the literature, [25, 26], apart from the above mentioned mass, one also defines the ADM mass in the Einstein frame, or tensor mass M_T , and the Keplerian mass, M_K . In the framework of a scalar tensor gravitational theory, these masses are obtained from the relations:

$$M_K = M + \Phi_1 \quad M_T = M + \frac{\Phi_1}{2} , \quad (31)$$

with

$$\Phi_1 = \lim_{r \rightarrow \infty} (r^2 \Phi'_{\text{BD}}) . \quad (32)$$

In our case we used the ADM mass in the Jordan frame for three reasons: It comes naturally from the Schwarzschild mass, we can deduce the Keplerian and tensor mass from eqs. 31 and 32 and in our case is always larger from M_K and M_T . For the $\omega_{\text{BD}} = -1$ case, and according to the literature, [25, 26], $\Phi'_{\text{BD}} > 0$ holds at infinity. For our case ($\omega_{\text{BD}} \geq 5$), we found numerically that $\Phi'_{\text{BD}} < 0$ not only at infinity but also for the whole region outside the soliton star. Because, in our case, $M > M_K, M_T$, and because the energy of the same number of free particles is larger than M , we find that the energy of the free particles is always larger than M_K and M_T , confirming the stability of the soliton with respect to fission into free particles.

The Noether current that leads to the conserved particle number is:

$$j^\mu = \imath \sqrt{-g_D} g^{\mu\nu} (\phi \partial_\nu \phi^* - \phi^* \partial_\nu \phi) , \quad (33)$$

which gives a conserved Noether charge:

$$N = \int_0^\infty d^{D-1} x j^t = \frac{4\pi^{(D-1)/2}}{\Gamma\left(\frac{D-1}{2}\right)} \int_0^R dr \omega \sigma^2 r^{D-2} \sqrt{\frac{B}{A}} . \quad (34)$$




Figure 6: The radius of a q-star as a function of the cosmological constant. In figures 6-10 dashed lines correspond to $\omega_{\text{BD}} = 500$ and solid lines to $\omega_{\text{BD}} = 5$.




Figure 7: The value of the matter scalar field at the center of a q-star as a function of the cosmological constant.




Figure 8: The mass of a q-star as a function of the cosmological constant.




Figure 9: The particle number of a q-star as a function of the cosmological constant.




Figure 10: The mass, M , and the particle number, N , of a q-star in $2 + 1$ dimensions as a function of the cosmological constant.

3 General scalar-tensor theory

In the original BD gravitational theory ω_{BD} is a constant. In a more general theory it may be regarded as a function, usually of the BD scalar. We use one of the simple functions, investigated in a cosmological framework, [33, 34], namely:

$$(D - 2)\omega_{\text{BD}} + D - 1 = \omega_0 \phi_{\text{BD}}^n, \quad (35)$$

with ω_0 and n constants. The equation of motion for the BD field is:

$$\phi_{\text{BD};\lambda}{}^{;\lambda} = \frac{1}{(D - 2)\omega_{\text{BD}} + D - 1} \left(8\pi T - \frac{d\omega_{\text{BD}}}{d\phi_{\text{BD}}} \phi_{\text{BD}}{}^{;\rho} \phi_{\text{BD};\rho} \right), \quad (36)$$

We set for simplicity $\Lambda = 0$. We rescale:

$$\tilde{\omega}_0 = \frac{(D - 2)\omega_{\text{BD}} + D - 1}{(D - 2)\omega_{\text{BD}} + D} G_D^m \omega_0, \quad (37)$$

and the other quantities as in eqs. 15-18. We will use for our calculations $n = 1$. For $n = 2, 3, 4$ the behavior of the star parameters is similar. Dropping




Figure 11: The radius of a q-star as a function of ω_0 .

the tildes and the $O(\epsilon)$ quantities we find for the Einstein equations:

$$G_t^t = \frac{\omega_0}{\omega_0 \Phi_{\text{BD}} + 1} \left[-W - U - \frac{1}{\omega_0 \Phi_{\text{BD}}} \times \right. \\ \left. \left((D-2)W - DU - \frac{A\Phi_{\text{BD}}'^2 \omega_0 \Phi_{\text{BD}} + 1}{D-2} \frac{1}{\Phi_{\text{BD}}} \right) \right] \\ - \frac{\omega_0 \Phi_{\text{BD}} - D + 1}{D-2} \frac{A\Phi_{\text{BD}}'^2}{2\Phi_{\text{BD}}^2} - \frac{AB'\Phi_{\text{BD}}'}{2\Phi_{\text{BD}}B}, \quad (38)$$

$$G_r^r = \frac{\omega_0}{\omega_0 \Phi_{\text{BD}} + 1} \left[W - U - \frac{1}{\omega_0 \Phi_{\text{BD}}} \times \right. \\ \left. \left((D-2)W - DU - \frac{A\Phi_{\text{BD}}'^2 \omega_0 \Phi_{\text{BD}} + 1}{D-2} \frac{1}{\Phi_{\text{BD}}} \right) \right] \\ + \frac{\omega_0 \Phi_{\text{BD}} - D + 1}{D-2} \frac{A\Phi_{\text{BD}}'^2}{2\Phi_{\text{BD}}^2} + \frac{A\Phi_{\text{BD}}''}{\Phi_{\text{BD}}} + \frac{A'\Phi_{\text{BD}}'}{2\Phi_{\text{BD}}}. \quad (39)$$

The equation of motion for the BD field is:

$$A \left[\Phi_{\text{BD}}'' + \left(\frac{D-2}{r} + \frac{A'}{2A} - \frac{B'}{2B} \right) \Phi_{\text{BD}}' \right] = \\ \frac{1}{\omega_0 \Phi_{\text{BD}} + 1} \left[(D-2)W - DU - \frac{A\Phi_{\text{BD}}'^2 \omega_0 \Phi_{\text{BD}} + 1}{D-2} \frac{1}{\Phi_{\text{BD}}} \right]. \quad (40)$$

We solve numerically the coupled system of eqs. 38-40. The mass and particle number of the star are given by eqs. 30, 34 respectively. We start




Figure 12: The value of the matter scalar field at the center of the q-star as a function of ω_0 .




Figure 13: The total energy of a q-star as a function of ω_0 .




Figure 14: The particle number of a q-star as a function of ω_0 .

from $\omega_0 = 5$ up to $\omega_0 = 1000$, where the results of general relativity are approximately reproduced.

4 Conclusions

In the present work we write the Brans-Dicke and a simple scalar-tensor gravitational theory in D dimensions and apply it in a special case of stable, gravitating, scalar field configuration, namely that of a q-star. The results of general relativity are reproduced for $\omega_{\text{BD}}, \omega_0 \rightarrow \infty$. A generally accepted, lower experimental limit is $\omega_{\text{BD}} \simeq 500$. Even for this value, the results almost coincide with General Relativity and one can regard that the results of General Relativity are practically obtained from our figures when $\omega_{\text{BD}}, \omega_0 = 1000$. We investigate the phase space from $\omega_{\text{BD}}, \omega_0 = 5$ up to 1000 so as to gain a detailed picture of the behavior of the star parameters, radius, value of the matter scalar field at the center of the star, total mass and particle number. For $\omega_{\text{BD}}, \omega_0 < 500$ the star parameters are a few per cent larger than in the case of General Relativity. These differences are D -dependent. For example, for $\omega_{\text{BD}} = 5$ the particle number is 33, 24, 21, 18 per cent larger from the corresponding results in General Relativity, for $D = 3, 4, 5$ and 6 dimensions respectively, the radius is 15, 7, 4 and 3 per cent larger respectively, and the total mass is $\sim 0, 7, 9$ and 10 per cent larger. One can find similar differences for the scalar-tensor theory.

The results of figures 1-5 and 11-14 refer to zero cosmological constant. Figure 5 shows the ratio M/N of the star as a function of its particle number

(M is the ADM star mass and N is the particle number). As one can see, every field configuration is stable with respect to fission into free particles. This feature is characteristic for solitonic gravitating objects, which are stable even in the absence of gravity. For example, a q-star with very small particle number and mass corresponds to a (non-gravitating) q-ball (in the thin-wall approximation). For small q-stars, one can see from figure 5 that $M/(mN) \rightarrow 0.5$, with m the mass of the free particles (here equal to unity with our rescalings). This is the correct limiting value in the absence of gravity, for the potential of 19, for which $\omega_{\min} \equiv (\sqrt{U/\sigma^2})_{\min} = 0.5$. The picture is absolutely different for non-solitonic boson stars, [35], where gravity and not the scalar potential is the main stabilizing factor. This means that for some region of the phase space, gravity is not strong enough to make the binding energy negative (equivalently, to make $M/(mN) < 1$). In figures 6-9 and for 4, 5 and 6 dimensions we start from a zero value of the cosmological constant, gradually decrease its value, and stop our calculations when $B(0) \rightarrow \infty$, or $\sigma(0) \rightarrow \infty$, showing in this way the formation of an anomaly at the center of the star. For $D = 3$, when decreasing the value of Λ we find that both mass and particle number increase up to their maximum value at approximately $\Lambda \simeq -3$, and below this value, the energy is approximately constant, when the particle number decreases slowly. Below $\Lambda \simeq -10$ the energy of the free particles (identified with the particle number for $m = 1$) is less than the soliton energy, which means that the decay into free particles is energetically favorable. All the other field configurations depicted in our figures are stable with respect to decay into free particles.

ACKNOWLEDGEMENTS

I wish to thank P. Manousselis and N. D. Tracas for helpful discussions.

References

- [1] D. J. Kaup, Phys. Rev. **172**, 1331 (1968).
- [2] R. Ruffini and S. Bonazzola, Phys. Rev. **187**, 1767 (1969).
- [3] E. W. Mielke and R. Scherzer, Phys. Rev. **D 24**, 2111 (1981).
- [4] M. Colpi, S. L. Shapiro, and I. Wasserman, Phys. Rev. Lett. **57**, 2485 (1986).

- [5] Ph. Jetzer and J. J. van der Bij, Phys. Lett. **B 227**, 341 (1989).
- [6] D. Astefanesei and E. Radu, Phys. Lett. **B 587**, 7 (2004).
- [7] R. Friedberg, T. D. Lee, and Y. Pang, Phys. Rev. **D 35**, 3640 (1987).
- [8] R. Friedberg, T. D. Lee, and Y. Pang, Phys. Rev. **D 35**, 3658 (1987).
- [9] R. Friedberg, T. D. Lee, and Y. Pang, Phys. Rev. **D 35**, 3678 (1987).
- [10] S. Coleman, Nucl. Phys. **B 262**, 263 (1985).
- [11] K. Lee, J. A. Stein-Schabes, R. Watkins, and L. M. Widrow, Phys. Rev. **D 39**, 1665 (1989).
- [12] A. M. Safian, S. Coleman, and M. Axenides, Nucl. Phys. **B 297**, 498 (1988).
- [13] B. W. Lynn, Nucl. Phys. **B 321**, 465 (1989).
- [14] S. Bahcall, B. W. Lynn, and S. B. Selipsky, Nucl. Phys. **B 325**, 606 (1989).
- [15] S. Bahcall, B. W. Lynn, and S. B. Selipsky, Nucl. Phys. **B 325**, 67 (1990).
- [16] A. Prikas, Phys. Rev. **D 66**, 025023 (2002).
- [17] A. Prikas, Gen. Rel. Grav. **36**, 1841 (2004).
- [18] A. Prikas, Phys. Rev. **D 69**, 125008 (2004).
- [19] C. Brans and R. H. Dicke, Phys. Rev. **124**, 925 (1961).
- [20] P. Jordan, Z. Phys. **157**, 112 (1959).
- [21] P. J. Bergmann, Int. J. Theor. Phys. **1**, 25 (1968).
- [22] R. V. Wagoner, Phys. Rev. **D 1**, 3209 (1970).
- [23] M. A. Gunderson and L. G. Jensen, Phys. Rev. **D 48**, 5628 (1993).
- [24] J. Balakrishna and Hisa-aki Shinkai, Phys. Rev. **D 58**, 044016 (1998).
- [25] A. W. Whinnett, Class. Quant. Grav. **16**, 2797 (1999).

- [26] D. F. Torres, F. E. Schunck and A. R. Liddle, *Class. Quant. Grav.* **15**, 3701 (1998).
- [27] F. E. Schunck and E. W. Mielke, *Class. Quant. Grav.* **20**, R301 (2003).
- [28] F. W. Hehl, J. D. McCrea, E. W. Mielke and Y. Ne'eman, *Phys. Rept* **258**, 1 (1995).
- [29] E. W. Mielke, *Phys. Rev.* **D 63**, 044018 (2001).
- [30] D. F. Torres, *Phys. Rev.* **D 56**, 3478 (1997).
- [31] G. L. Comer and Hisa-aki Shinkai, *Class. Quant. Grav.* **15**, 669 (1998).
- [32] A. W. Whinnet and D. F. Torres, *Phys. Rev.* **D 60**, 104050 (1999).
- [33] J. D. Barrow and P. Parsons, *Phys. Rev.* **D 55**, 1906 (1997).
- [34] J. D. Barrow and J. P. Mimoso, *Phys. Rev.* **D 50**, 3746 (1994).
- [35] F. V. Kusmartsev, E. W. Mielke and F. E. Schunck, *Phys. Rev.* **D 43**, 3895 (1991).